

## **3-D Visualization of Morphology and Distribution of Acoustic Facies in the Strataform Study Areas**

Larry Mayer  
Ocean Mapping Group  
Dept. of Geodesy and Geomatics Engineering  
University of New Brunswick  
Fredericton, N.B. Canada E3B 5A3  
phone: (506) 453-3577 fax: (506) 453-4943 email: [larry@omg.unb.ca](mailto:larry@omg.unb.ca)  
Award # N00014-95-0064  
<http://www.omg.unb.ca>

### **LONG-TERM GOAL**

Our long-term goal is to provide the fundamental mapping information and visualization tools necessary to support STRATAFORM's multidisciplinary effort to develop a more complete understanding of how short-term oceanographic and geological processes interact to produce the preserved geologic record on the shelf and slope portions of the continental margins.

### **SCIENTIFIC OBJECTIVES**

The fundamental objective of the STRATAFORM Swathmapping Program is **to provide complete (100%) bathymetric and sidescan imagery coverage of the Northern California and N.J. Margin STRATAFORM field areas**. This has allowed STRATAFORM investigators to evaluate the geologic processes of the shelf and slope over a continuum of scales. Complete coverage also has provided STRATAFORM investigators with the knowledge that their studies are based on a complete picture of morphological relationships rather than the interpolation of sparsely spaced data. In doing this, we have produced a bathymetric, geomorphological, and potentially lithological framework upon which all subsequent work can be built. Ancillary objectives are **to further develop techniques for the remote classification of seafloor materials from swathmapping data and, to develop techniques for the interactive 3D visualization of co-registered surficial and subbottom data**.

### **APPROACH**

We have chosen to use a state-of-the-art multibeam sonar system (EM-1000) for our mapping. As compared to conventional echosounders, multibeam sonars provide increased source level, lateral resolution, and a substantial increase in data density and areal coverage. Most importantly, the newer systems also provide the ability to simultaneously produce high-resolution sidescan sonar imagery. We have developed a full suite of real-time and near real-time multibeam sonar processing tools to assure that only high-quality data is collected and that this data can be processed in the field. These tools also allow for the interactive 3-D visualization of multibeam data sets and derivative products. We are also developing a range of seafloor classification algorithms with particular focus on techniques that look at the characteristics of the returned waveforms as well as the change in backscatter as a function of angle of incidence. We are also exploring the application of the wavelet transform as a means of removing

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>3-D Visualization of Morphology and Distribution of Acoustic Facies in the Strataform Study Areas</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of New Brunswick, Department of Geodesy and Geomatics Engineering, Ocean Mapping Group, Fredericton, N.B. Canada E3B 5A3,</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

unwanted artifacts from seafloor backscatter data as well as a means of estimating seafloor type. The results of these analyses will be compared to ground truth data collected by other STRATAFORM scientists to explore the limits of extracting quantitative seafloor property information from multibeam sonar data.


## **WORK COMPLETED**

In 1995 and 1996, we completed mapping of the Calif. and most of the N.J. survey areas. In 1997 we completed the remaining deep-water portion of the N.J., processed these data and merged them with the shallow water data. We also processed additional multibeam (Hydrosweep) data from the Eureka area collected by Clark Alexander. We merged this data with the earlier data sets and made the new maps available to all STRATAFORM investigators. We have now focused our efforts on visualization and particularly on the question of remote seafloor classification. We have finalized interactive 3-D fly-throughs of the N. J. margin and have developed a suite of automated algorithms for extracting and parameterizing the backscatter as a function of angle of incidence. We have also nearly completed the software necessary to bring fully georeferenced high-resolution seismic data into our 3-D visualization package. Our focus for sediment classification studies has been the Eureka margin. Seismic data, gas abundance data, as well as data from box cores, piston cores, and camera stations have all been entered into a 2-D GIS system for direct comparison with backscatter data. Work is underway to transition these data sets into our interactive 3-D GIS. Finally, we have completed the algorithms for the application of tree-structured wavelet transforms to multibeam sonar backscatter data and applied this transform to data from the Eureka margin.


## **RESULTS**

The combined bathymetry and imagery of the N. J. margin show a spectacular picture of shelf processes with large scale ridge and swale features that appear to be moving across the shelf, a series of ablation features and, a large region near the shelf break that appears to be covered by old iceberg scours. The ridge and swale topography appears to be overriding the scour features though there are several large scour-like feature that also seem to cut into the ridge and swale deposits. If these large features are iceberg scours, this part of the system has been preserved since the last deglaciation. The deep water data provides direct insight into the origin of submarine canyons, with landward erosion apparently the dominant mechanism. Our parameterization of the angular backscatter response curve appears to be a sensitive indicator of changes in seafloor type or texture. In particular, the shape of the curve below  $40^\circ$  is especially revealing as response in the vicinity of the critical angle may be observed. The use of our 3-D visualization tools has allowed us to directly see the spatial distribution of our extracted classifiers and their relationship to seafloor morphology.

The wavelet transform has proven to be a robust technique for the removal of artifacts from backscatter data. Backscatter data characteristically is degraded by ship-track parallel artifacts that are the result of the incomplete correction for specular energy at nadir (i.e., the lack of sidescan at nadir). These artifacts are easily identified as “channels” at certain levels of the wavelet transform and thus easily removed (Fig 1). Once the artifacts are removed we can continue to use the wavelet transform to identify regions of characteristic backscatter statistics that we then relate to changes in seafloor roughness. We are using quantitative roughness data provided by other Strataform investigators to ground truth these studies.



***Fig 1. A. EM-1000 backscatter data with dominant ship-track parallel nadir artifacts***



***Fig 1 B. Backscatter data from Fig 1A with artifact removed via wavelet transform***

## **IMPACT/APPLICATIONS**

The swathmapping results from the Eureka and the N. J. margins provide all STRATAFORM investigators with an unprecedented, detailed look at both the bathymetry and distribution of sediments on the shelf and slope. These results have already been used in planning the deployment of a series of long-term moorings, seismic profiling and coring cruises, as well as for planning ROV, submersible work, and ODP sites. Quantitative measurements of slope and sediment-type distributions will inevitably lead to improved models of shelf and slope development, the primary goal of the STRATAFORM program. Our initial results from the New Jersey margin indicate that pervasive ice scouring is evident further south than previously reported and may have impact on our overall understanding of the evolution of shelf stratigraphy. The ability to interactively explore large and complex data sets in a 3-D GIS will greatly facilitate our understanding of the seafloor processes at work in the Strataform areas and particularly their relationship to seafloor morphology. In addition, through the use of backscatter as a function of angle of incidence and wavelet transforms we hope to provide a new and rapid means of exploring lateral changes in seafloor roughness and composition.

## **TRANSITIONS**

Our maps and data have been used by numerous investigators both in and out of STRATAFORM, including several other Navy programs (e.g., SWARM, NRL). Our processing and visualization software is being used by NAVO, NRL, NOAA, USGS as well as a number of universities and private sector companies (Shell, Exxon, Woodside, WAPET).

## **RELATED PROJECTS**

NSERC Chair in Ocean Mapping – a consortium of 14 sponsors that provide base support for general research in the field of ocean mapping.

## **PUBLICATIONS**

Mayer, L.A., Li, Yanchao, Melvin, G., and Ware, C., The application of 3-D visualization technology to pelagic fisheries assessment and research, ICES, in press.

Gardner, J.M, Mayer, L.A., Hughes Clarke, J.E.H., and Kleiner, A., Quantitative bathymetry of East Flower Gardens and Stetson Bank, Gulf of Mexico, *Journal of Gulf of Mexico Science*, in press

Goff, J., Orange, D.L., Mayer, L.A., and Hughes Clarke, J.E., Detailed investigation of continental shelf morphology from a high-resolution swath sonar survey over the Eel River Basin, Northern California, *Marine Geology*, in press

Borgeld, J.C., Hughes Clarke, J.E., Goff, J.A., Mayer, L.A., and Curtis, J.A., Acoustic backscatter of the 1995 flood deposit on the Eel Shelf, *Marine Geology*. in press.

Brew, D. and Mayer, L.A., 1998, Modelling of Pliocene-Pleistocene abyssal mudwaves using synthetic seismograms: *Marine Geol.*, v. 149, pp. 3-16.

Gardner, J. V., Butman, P.B., Mayer, L.A. and Hughes Clarke, J.E., 1998, Mapping U.S. continental shelves, *Sea Technology*, v. 39., no. 6, pg. 10 – 18.

Hughes Clarke, J.E., 1998, The effect of fine scale seabed morphology and texture on the fidelity of swath bathymetric sounding data, *Proceedings, Canadian Hydrographic Conference*, April 1998, vol. 1, pg. 168 – 181.

Hughes Clarke, J.E., Gardner, J.V., Torreson, M., and Mayer, L.A., 1998, The limits of spatial resolution achievable using a 30 kHz multibeam sonar: model predictions and field results., in *IEEE Oceans 98, Conference Proceedings*, v. 3. pg. 1823 – 1827.

Paton, M., Mayer, L.A., and Ware, C., 1997, Interactive 3-D tools for pipeline route planning, *Proceedings of the IEEE- Oceans 97*, Vol. 2, pg. 1216 - 1222

Mayer, L.A., Dijkstra, S., Hughes Clarke, J.E., Paton, M. and Ware, C., 1997, Interactive tools for the exploration and analysis of multibeam and other seafloor acoustic data, in Pace, N.G., Pouliquen, E., Bergem, O., and Lyons., eds., *High Frequency Acoustics in Shallow Water*, SACLANT Conference Proceedings Series, CP-45, NATO SACLANT Research Centre, La Spezia, Italy, p. 355 - 362.

Brissette, M.B., Hughes Clarke, J.E., Bradford, J. and MacGowan, B., 1997, Detecting small seabed targets using a high frequency multibeam sonar: *Geometric Models and Test Results: Proc. Oceans 97*.